

# BREO ELLIPTA

**The only once-daily ICS/LABA  
(inhaled corticosteroid/long-acting beta<sub>2</sub>-agonist)  
for the maintenance treatment of COPD  
(chronic obstructive pulmonary disease).**



## Indications

- BREO ELLIPTA is a combination inhaled corticosteroid/long-acting beta<sub>2</sub>-adrenergic agonist (ICS/LABA) indicated for the long-term, once-daily, maintenance treatment of airflow obstruction in patients with chronic obstructive pulmonary disease (COPD), including chronic bronchitis and/or emphysema. BREO ELLIPTA is also indicated to reduce exacerbations of COPD in patients with a history of exacerbations.
- BREO ELLIPTA is NOT indicated for the relief of acute bronchospasm or for the treatment of asthma.

## Important Safety Information for BREO ELLIPTA

### WARNING: ASTHMA-RELATED DEATH

- Long-acting beta<sub>2</sub>-adrenergic agonists (LABAs), such as vilanterol, one of the active ingredients in BREO ELLIPTA, increase the risk of asthma-related death. A placebo-controlled trial with another LABA (salmeterol) showed an increase in asthma-related deaths in subjects receiving salmeterol. This finding with salmeterol is considered a class effect of all LABAs, including vilanterol.
- The safety and efficacy of BREO ELLIPTA in patients with asthma have not been established. BREO ELLIPTA is not indicated for the treatment of asthma.

## CONTRAINDICATIONS

- BREO ELLIPTA is contraindicated in patients with severe hypersensitivity to milk proteins or who have demonstrated hypersensitivity to either fluticasone furoate, vilanterol, or any of the excipients.

## WARNINGS AND PRECAUTIONS

- BREO ELLIPTA should not be initiated in patients during rapidly deteriorating or potentially life-threatening episodes of COPD.
- BREO ELLIPTA should not be used for the relief of acute symptoms, i.e., as rescue therapy for the treatment of acute episodes of bronchospasm. Acute symptoms should be treated with an inhaled, short-acting beta<sub>2</sub>-agonist.
- BREO ELLIPTA should not be used more often than recommended, at higher doses than recommended, or in conjunction with other medications containing LABAs, as an overdose may result. Clinically significant cardiovascular effects and fatalities have been reported in association with excessive use of inhaled sympathomimetic drugs. Patients using BREO ELLIPTA should not use another medicine containing a LABA (e.g., salmeterol, formoterol fumarate, arformoterol tartrate, indacaterol) for any reason.
- Oropharyngeal candidiasis has occurred in patients treated with BREO ELLIPTA. Advise patients to rinse the mouth without swallowing following inhalation to help reduce the risk of oropharyngeal candidiasis.
- An increase in the incidence of pneumonia has been observed in subjects with COPD receiving BREO ELLIPTA. There was also an increased incidence of pneumonias resulting in hospitalization. In some incidences these pneumonia events were fatal.
  - In replicate 12-month studies of 3255 subjects with COPD who had experienced a COPD exacerbation in the previous year, there was a higher incidence of pneumonia reported in subjects receiving BREO ELLIPTA 100/25 mcg (6% [51 of 806 subjects]), fluticasone furoate (FF)/vilanterol (VI) 50/25 mcg (6% [48 of 820 subjects]), and FF/VI 200/25 mcg (7% [55 of 811 subjects]) than in subjects receiving VI 25 mcg (3% [27 of 818 subjects]). There was no fatal pneumonia in subjects receiving VI or FF/VI 50/25 mcg. There was fatal pneumonia in 1 subject receiving BREO ELLIPTA at the approved strength (100/25 mcg) and in 7 subjects receiving FF/VI 200/25 mcg (<1% for each treatment group).
- Physicians should remain vigilant for the possible development of pneumonia in patients with COPD, as the clinical features of such infections overlap with the symptoms of COPD exacerbations.
- Patients who use corticosteroids are at risk for potential worsening of existing tuberculosis; fungal, bacterial, viral, or parasitic infections; or ocular herpes simplex. A more serious or even fatal course of chickenpox or measles may occur in susceptible patients. Use caution in patients with the above because of the potential for worsening of these infections.
- Particular care is needed for patients who have been transferred from systemically active corticosteroids to inhaled corticosteroids because deaths due to adrenal insufficiency have occurred in patients with asthma during and after transfer from systemic corticosteroids to less systemically available inhaled corticosteroids. Taper patients slowly from systemic corticosteroids if transferring to BREO ELLIPTA.

Please see additional Important Safety Information for BREO ELLIPTA on back.

Please see accompanying full Prescribing Information, including Boxed Warning and Medication Guide, for BREO ELLIPTA.

